

Briarane Diterpenoids from the Formosan Gorgonian Coral *Junceella fragilis*

Ping-Jyun SUNG,* Mei-Ru LIN, and Lee-Shing FANG

National Museum of Marine Biology and Aquarium, 2 Houwan Road, Checheng, Pingtung 944, Taiwan, R.O.C.

Received July 28, 2004; accepted September 3, 2004

A new trihydroxy briarane-related diterpenoid, junceollide I (1), along with a known metabolite, (1R,2R,5Z,7R,8S,9R,10R,12R,14R,17S)-2,14-diacetoxy-8,17-epoxy-9,12-dihydroxybriara-5,11(20)-dien-19-one (2), have been isolated from the gorgonian coral *Junceella fragilis*, collected off the southern Taiwan coast. The structure, including the relative configuration of the new compound 1, was elucidated by the combination of spectral data, particularly in high-resolution ¹H- and ¹³C-NMR spectroscopy utilizing COSY, HMBC, HMQC, and NOESY experiments.

Key words *Junceella fragilis*; junceollide; gorgonian; briarane

Previous chemical investigations of the gorgonian coral *Junceella fragilis* (phylum Cnidaria, class Anthozoa, subclass Octocorallia, order Gorgonacea, family Ellisellidae)^{1–3} have yielded 15 new diterpenoids possessing the briarane skeleton. These metabolites are junceollides A—H,^{4–6} (–)-4-deacetyljunceollide D, (+)-11 α ,20 α -epoxyjunceollide D, (–)-11 α ,20 α -epoxy-4-deacetyljunceollide D, (–)-11 α ,20 α -epoxy-4-deacetoxyjunceollide D, (+)-junceollide A,⁷ 9-O-deacetylumbraculolide A,⁸ and fragilide A.⁹ In addition to species of the genus *Junceella*,¹⁰ briarane-type metabolites have also been isolated from a variety of marine organisms, and the compounds of this type were found to possess extensive biological activity,¹¹ and could be originally synthesized by the corals.^{12,13} In this paper, we report the isolation and structure determination of two briarane derivatives, including a new briarane, junceollide I (1), together with a known metabolite, (1R,2R,5Z,7R,8S,9R,10R,12R,14R,17S)-2,14-diacetoxy-8,17-epoxy-9,12-dihydroxybriara-5,11(20)-dien-19-one (2),¹⁴ from the gorgonian coral *Junceella fragilis* collected off southern Taiwan coast. The structure of the new diterpenoid 1 was elucidated by combined analysis of spectral data.

Junceollide I (1) was obtained as white powder, [α]_D²⁵ –77° (*c*=0.7, CHCl₃). This metabolite has a molecular formula of C₂₄H₃₆O₉, as established by FAB-MS and NMR data, which indicates seven degrees of unsaturation. The IR absorptions of 1 showed the presence of hydroxy (3352 cm⁻¹), γ -lactone (1775 cm⁻¹), and ester (1736 cm⁻¹) groups. The FAB-MS of 1 exhibited peaks at *m/z* 451 (M+H–H₂O)⁺, 433 (M+H–2H₂O)⁺, 391 (M+H–AcOH–H₂O)⁺, 373 (M+H–AcOH–2H₂O)⁺, 349 (M+H–2AcOH)⁺, 331 (M+H–2AcOH–H₂O)⁺, 313 (M+H–2AcOH–2H₂O)⁺, and 295 (M+H–2AcOH–3H₂O)⁺, also indicating the presence of two acetoxy and three hydroxy groups in 1. From the ¹³C-NMR data of 1 (Table 1), the presence of a trisubstituted olefin was deduced from the signals of two carbons resonating at δ _C 146.2 (s) and 119.0 (d). Furthermore, in the ¹³C-NMR spectrum, three carbonyl resonances appeared at δ _C 176.2 (s), 170.3 (s), and 169.4 (s), confirming the presence of a γ -lactone and two esters in 1. In the ¹H-NMR spectrum of 1 (Table 1), two acetyl methyl groups (δ _H 2.18, 3H, s; 2.06, 3H, s) were observed. Thus, the NMR data accounted for four degrees of unsaturation and requiring 1 to be tricyclic.

The ¹H-NMR spectrum also showed the presence of four methyl groups, including a methyl (δ _H 1.16, 3H, d, *J*=7.5 Hz, H₃-18) attached to a methine carbon, a methyl (δ _H 0.97, 3H, s, H₃-15) attached to a tertiary carbon, a methyl (δ _H 1.43, 3H, s, H₃-20) attached to an oxygen-bearing quaternary carbon, and a vinyl methyl (δ _H 2.03, 3H, s, H₃-16).


The gross structure of 1 and all of the ¹H- and ¹³C-NMR data associated with the molecule were determined by 2D NMR studies, including ¹H–¹H correlation spectroscopy (COSY), ¹H-detected heteronuclear multiple quantum coherence (HMQC), and heteronuclear multiple bond connectivity (HMBC) experiments. From the ¹H–¹H COSY spectrum of 1 (Fig. 1), it was possible to establish the separate spin systems that map out the proton sequences from H-2/H₂-3; H₂-3/H₂-4; H-6/H-7; and H-9/H-10. These data, together with the HMBC correlations between H-2/C-1, C-3, C-4, C-10; H₂-3/C-1, C-2, C-4, C-5; H₂-4/C-3, C-5, C-6; H-7/C-5, C-6, C-8; H-9/C-7, C-8, C-10; and H-10/C-1, C-2, C-8, C-9 (Fig. 1, Table 1), established the connectivity from C-1 to C-10 within the ten-membered ring. A vinyl methyl attached at C-5 was confirmed by the long-range ¹H–¹H COSY correlations between H₃-16 and H-6 and the HMBC correlations between H₃-16/C-4, C-5, and C-6. The methylcyclohexane ring was elucidated by the combination of ¹H–¹H COSY correlations between H₂-12/H₂-13 and H₂-13/H-14 and the HMBC correlations between H₂-12/C-11, C-20; H₂-13/C-1, C-11, C-12, C-14; H-14/C-10, C-12, C-13; and H₃-20/C-10, C-11, C-12. The methylcyclohexane ring, which is fused to the ten-membered ring at C-1 and C-10, was elucidated by the long-range *W*-coupling between H-10 and H-12 α and by the key HMBC correlations between H-2/C-14; H-9/C-11; H-10/C-11; and H-14/C-2. The ring-juncture C-15 methyl group was positioned at C-1 from the key HMBC correlations between H₃-15/C-1, C-2, C-10, C-14. Furthermore, the acetoxy groups positioned at C-2 and C-9 were confirmed from the HMBC correlations between δ _H 4.51 (H-2), 5.18 (H-9) and the ester carbonyl carbons appeared at δ _C 170.3 (s), 169.4 (s), respectively. In addition, the proton of the hydroxy-bearing methine showed the signal at δ _H 4.12 (1H, d, *J*=5.0 Hz) was assigned to H-14. The 11-hydroxy group was confirmed from the signal of a quaternary oxygen-bearing carbon at δ _C 89.1 (s) and from the chemical shift of the tertiary methyl H₃-20 (δ _H 1.43, 3H, s). Thus, the remaining

* To whom correspondence should be addressed. e-mail: pjsung@nmmba.gov.tw

Table 1. ^1H - and ^{13}C -NMR Chemical Shifts and HMBC and ^1H - ^1H COSY Correlations for **1**


C/H	$^1\text{H}^a$	$^{13}\text{C}^b$	HMBC (H \rightarrow C)	^1H - ^1H COSY
1		51.6 (s) ^d	H-2, H ₂ -3, H-10, H ₂ -13, H ₃ -15	
2	4.51 (1H, t, $J=5.0$ Hz) ^c	77.4 (d)	H ₂ -3, H-4 α , H-10, H-14, H ₃ -15	H-3 α/β
3 α	1.91 m	32.6 (t)	H-2, H ₂ -4	H-2, H-3 β , H-4 α/β
β	2.04 m			H-2, H-3 α , H-4 α/β
4 α	2.29 (1H, ddd, $J=15.0, 10.5, 5.0$ Hz)	28.9 (t)	H-2, H ₂ -3, H ₃ -16	H-3 α/β ; H-4 β
β	2.59 (1H, dt, $J=15.0, 5.0$ Hz)			H-3 α/β ; H-4 α
5		146.2 (s)	H ₂ -3, H ₂ -4, H-7, H ₃ -16	
6	5.46 (1H, d, $J=9.5$ Hz)	119.0 (d)	H ₂ -4, H-7, H ₃ -16	H-7, H ₃ -16
7	5.27 (1H, d, $J=9.5$ Hz)	77.6 (d)	H-9, OH-8	H-6
8		80.9 (s)	H-7, H-9, H-10, H-17, H ₃ -18, OH-8	
9	5.18 (1H, d, $J=6.0$ Hz)	68.3 (d)	H-10, H-17, OH-8	H-10
10	2.13 (1H, dd, $J=6.0, 2.0$ Hz)	49.9 (d)	H-2, H-9, H ₂ -12, H-14, H ₃ -15, H ₃ -20	H-9, H-12 α
11		89.1 (s)	H-9, H-10, H ₂ -12, H ₂ -13, H ₃ -20	
12 α	1.30 (1H, m)	29.2 (t)	H-10, H ₂ -13, H-14, H ₃ -20	H-10, H-12 β , H-13 α/β
β	2.00 (1H, m)			H-12 α , H-13 α/β
13 α	2.88 (1H, m)	27.9 (t)	H-14	H-12 α/β , H-13 β , H-14
β	1.68 (1H, m)			H-12 α/β , H-13 α , H-14
14	4.12 (1H, d, $J=5.0$ Hz)	82.1 (d)	H-2, H ₂ -13, H ₃ -15	H-13 α/β
15	0.97 (3H, s)	15.4 (q)	H-2, H-10	
16	2.03 (3H, s)	26.5 (q)	H-4 α , H-6	H-6
17	2.43 (1H, q, $J=7.5$ Hz)	42.0 (d)	H-9, H ₃ -18, OH-8	H ₃ -18
18	1.16 (3H, d, $J=7.5$ Hz)	6.6 (q)	H-17	H-17
19		176.2 (s)	H-17, H ₃ -18	
20	1.43 (3H, s)	23.2 (q)	H-10, H ₂ -12	
OH-8	1.97 (1H, s)			
Acetates	2.18 (3H, s)	21.4 (q)		
		169.4 (s)	H-9, acetate methyl	
	2.06 (3H, s)	21.1 (q)		
		170.3 (s)	H-2, acetate methyl	

^a Measured at 500 MHz in CDCl_3 at 25 °C. ^b Measured at 125 MHz in CDCl_3 at 25 °C. ^c J values (in Hz) in parentheses. ^d Multiplicity deduced by DEPT and indicated by usual symbols.

Fig. 1. The ^1H - ^1H COSY and HMBC Correlations of **1**

hydroxy group had to be positioned at C-8, an oxygen-bearing quaternary carbon resonating at δ_{C} 80.9. The latter was further confirmed from the HMBC correlations observed between OH-8/C-7, C-8, C-9, C-17. These data, together with the HMBC correlations between H₃-18/C-8, C-17, C-19, unambiguously established the molecular framework of **1**.

The relative stereochemistry of **1** was elucidated from the NOE interactions observed in a nuclear overhauser effect spectroscopy (NOESY) experiment (Fig. 2). In the NOESY experiment of **1**, H-10 gives NOE correlations to H-2, OH-8,

Fig. 2. Selective NOE Correlations of **1**

and H₃-20, but not to H₃-15, and H-2 was found to show NOE responses with one proton of the C-3 methylene (δ_{H} 1.91, m, H-3 α), indicating that these protons are located on the same face of the molecule and assigned as α -protons, since C-15 methyl and 11-hydroxy groups are the β -substituents at C-1 and C-11, respectively, and 8-hydroxy group was α -oriented. H-14 was found to exhibit NOE responses with H₃-15, but not with H-10, revealing the β -orientation of this proton. Furthermore, H-7 was found to exhibit NOE correlations with H-3 β , one proton of the C-4 methylene (δ_{H} 2.59, 1H, dt, $J=15.0, 5.0$ Hz, H-4 β), and H-17, but not with OH-8, indicating H-7 and H-17 should be placed on the β face and H₃-18 was α -oriented in **1**. H-9 was found to show NOE responses with H₃-18 and H₃-20. From the consideration of molecular models, H-9 was found to be reasonably close to H₃-18 and H₃-20, thus being concluded that H-9 should be placed on the α face in **1**. From the above results, the structure, including the relative configuration of **1**, was established unambiguously.

Compound **2** was identified as a known diterpene,

(1*R*,2*R*,5*Z*,7*R*,8*S*,9*R*,10*R*,12*R*,14*R*,17*S*)-2,14-diacetoxy-8,17-epoxy-9,12-dihydroxybriara-5,11(20)-dien-19-one, which had been isolated from an Australian gorgonian coral *Junceella gemmacea*. Its physical (optical rotation value) and spectral (¹H- and ¹³C-NMR) data are in full agreement with those reported previously.¹⁴⁾

Experimental

Melting points were determined using a FARGO apparatus and were uncorrected. Optical rotation values were measured in CHCl₃ using a JASCO D-370 digital polarimeter at 25 °C. Infrared spectra were recorded on a JASCO 5300 FT-IR. FAB-MS was obtained with a VG QUATTRO GC/MS spectrometer. HR-FAB-MS was recorded on a VG 70-250S GC/MS spectrometer. NMR spectra were recorded a VARIAN UNITY INOVA 500 FT-NMR at 500 MHz for ¹H and 125 MHz for ¹³C, respectively, in CDCl₃ using TMS as an internal standard. Silica gel (Merck, 230-400 mesh) was used for column chromatography. TLC spots (Si gel 60 F₂₅₄, 0.2 mm, Merck) were detected with an UV₂₅₄ lamp and by 20% H₂SO₄ followed by heating at 120 °C for 5 min. All solvents used were either freshly distilled or of analytical grade.

Animal Material The gorgonian coral *Junceella fragilis* was collected by hand using scuba gear off the southern Taiwan coast on Dec. 12, 2002, at a depth of -10 m. Taxonomic identification was provided by Dr. Tung-Yung Fan from the National Museum of Marine Biology and Aquarium (NMMBA), R.O.C. The living reference specimen was deposited in the NMMBA (TWGC-003). This organism was identified from descriptions.¹⁻³⁾

Extraction and Isolation The organism (780 g) was collected and freeze-dried. The freeze-dried material (557 g) was minced and extracted with EtOAc (5×500 ml) for 120 h at 25 °C. The organic extract (11.1 g) was separated by silica gel column chromatography using *n*-hexane and *n*-hexane-EtOAc mixtures of increasing polarity. Briarane **2** was eluted with *n*-hexane-EtOAc (5 : 2) and **1** with *n*-hexane-EtOAc (1 : 1).

Junceollolide I (**1**): White powder (7.8 mg); mp 210-212 °C; [α]_D²⁵ -77° (*c*=0.7, CHCl₃); IR (neat) cm⁻¹ 3352, 1775, 1736; ¹H- and ¹³C-NMR data, see Table 1; FAB-MS *m/z*: 451, 433, 391, 373, 349, 331, 313, 295. HR-FAB-MS: *m/z* 451.2332 (Calcd for C₂₄H₃₆O₉+H-H₂O, 451.2333).

(1*R*,2*R*,5*Z*,7*R*,8*S*,9*R*,10*R*,12*R*,14*R*,17*S*)-2,14-diacetoxy-8,17-epoxy-9,12-dihydroxybriara-5,11(20)-dien-19-one (**2**): Colorless oil (1.0 mg); [α]_D²⁵ +113° (*c*=0.1, CHCl₃) (lit.¹⁴⁾ [α]_D +115.1° (*c*=0.08)); ¹H-NMR (500 MHz, CDCl₃) δ 5.53 (1H, d, *J*=9.0 Hz), 5.34 (1H, d, *J*=9.0 Hz), 5.31 (1H, s), 5.14 (1H, d, *J*=7.0 Hz), 5.04 (1H, s), 4.77 (1H, brs), 4.35 (1H, brs), 4.31 (1H, t,

J=7.0 Hz), 3.04 (1H, brs), 2.65 (1H, brt, *J*=15.5 Hz), 2.57 (1H, brd, *J*=15.5 Hz), 2.18 (1H, m), 2.00 (3H, s), 1.98 (3H, s), 1.96 (3H, s), 1.88 (1H, m), 1.78 (1H, m), 1.70 (1H, m), 1.53 (3H, s), 1.26 (3H, s); ¹³C-NMR (125 MHz, CDCl₃) δ 171.8 (s), 170.6 (s), 170.2 (s), 151.7 (s), 144.5 (s), 119.3 (d), 110.5 (t), 75.0 (d), 74.3 (d), 74.2 (d), 73.6 (d), 71.4 (s), 69.7 (d), 62.2 (s), 47.2 (s), 44.0 (d), 36.4 (t), 31.2 (t), 28.5 (t), 26.9 (q), 21.4 (q), 21.1 (q), 15.0 (q), 10.0 (q). The related physical (optical rotation value) and spectral (¹H- and ¹³C-NMR) data of **2** are in full agreement with those reported previously.¹⁴⁾

Acknowledgments This work was supported by the grants from the National Museum of Marine Biology and Aquarium and the National Science Council (Contract no. NSC 92-2323-B-291-001 and 92-2320-B-291-001) of the Republic of China awarded to P.-J. Sung. We thank Dr. Tung-Yung Fan, the National Museum of Marine Biology and Aquarium, R.O.C., for his collection and identification of the marine organisms.

References

- 1) Bayer F. M., *Proc. Biol. Soc. Wash.*, **94**, 902-947 (1981).
- 2) Chen C.-C., Chang K.-H., *Bull. Inst. Zool., Academia Sinica*, **30**, 149-182 (1991).
- 3) Bayer F. M., Grasshoff M., *Senckenbergiana Biol.*, **74**, 21-45 (1994).
- 4) Shin J., Park M., Fenical W., *Tetrahedron*, **45**, 1633-1638 (1989).
- 5) Sung P.-J., Wu S.-L., Fang H.-J., Chiang M. Y., Wu J.-Y., Fang L.-S., Sheu J.-H., *J. Nat. Prod.*, **63**, 1483-1487 (2000).
- 6) Sung P.-J., Fan T.-Y., Fang L.-S., Wu S.-L., Li J.-J., Chen M.-C., Cheng Y.-M., Wang G.-H., *Chem. Pharm. Bull.*, **51**, 1429-1431 (2003).
- 7) García M., Rodríguez J., Jiménez C., *J. Nat. Prod.*, **62**, 257-260 (1999).
- 8) Sung P.-J., Fan T.-Y., *Heterocycles*, **60**, 1199-1202 (2003).
- 9) Sung P.-J., Lin M.-R., Chen W.-C., Fang L.-S., Lu C.-K., Sheu J.-H., *Bull. Chem. Soc. Jpn.*, **77**, 1229-1230 (2004).
- 10) Sung P.-J., Gwo H.-H., Fan T.-Y., Li J.-J., Dong J., Han C.-C., Wu S.-L., Fang L.-S., *Biochem. Syst. Ecol.*, **32**, 185-196 (2004).
- 11) Sung P.-J., Sheu J.-H., Xu J.-P., *Heterocycles*, **57**, 535-579 (2002), and related references cited therein.
- 12) Kokke W. C. M. C., Epstein S., Look S. A., Rau G. H., Fenical W., Djerassi C., *J. Biol. Chem.*, **259**, 8168-8173 (1984).
- 13) Sung P.-J., Fan T.-Y., Chen M.-C., Fang L.-S., Lin M.-R., Chang P.-C., *Biochem. Syst. Ecol.*, **32**, 111-113 (2004).
- 14) Bowden B. F., Coll J. C., König G. M., *Aust. J. Chem.*, **43**, 151-159 (1990).